

Constraining a possible time-variation of the gravitational constant through “gravitochemical heating” of neutron stars

Paula Jofré,¹ Andreas Reisenegger,¹ and Rodrigo Fernández²

¹*Departamento de Astronomía y Astrofísica, Pontificia Universidad Católica de Chile, Casilla 306, Santiago 22, Chile*

²*Department of Astronomy & Astrophysics, University of Toronto, Toronto, ON M5S 3H8, Canada.*

(Dated: February 5, 2008)

A hypothetical time-variation of the gravitational constant G would cause neutron star matter to depart from beta equilibrium, due to the changing hydrostatic equilibrium. This forces non-equilibrium beta processes to occur, which release energy that is invested partly in neutrino emission and partly in heating the stellar interior. Eventually, the star arrives at a stationary state in which the temperature remains nearly constant, as the forcing through the change of G is balanced by the ongoing reactions. Comparing the surface temperature of the nearest millisecond pulsar, PSR J0437-4715, inferred from ultraviolet observations, with our predicted stationary temperature, we estimate two upper limits for this variation: (1) $|\dot{G}/G| < 2 \times 10^{-10} \text{ yr}^{-1}$, if we allow direct Urca reactions operating in the neutron star core, and (2) $|\dot{G}/G| < 4 \times 10^{-12} \text{ yr}^{-1}$, considering only modified Urca reactions. Both results are competitive with those obtained by other methods, with (2) being among the most restrictive.

PACS numbers: 91.10.Op, 06.20.Jr, 97.60.Jd

Since Dirac [1] suggested that the gravitational force may be weakening with the expansion of the universe, the question about whether or not the fundamental constants of nature vary in time has been of interest. General relativity assumes a strictly constant gravitational coupling parameter G , but alternatives such as the Brans-Dicke scalar-tensor theory [2] predict a variable G .

There have been many experiments attempting to test the constancy of G , obtaining upper limits for its variation, most usefully expressed as $|\dot{G}/G|$. The experiments can be separated into three classes. In the first class are those that measure the variation of G from the early Universe to the present time, over timescales $\gtrsim 10^{10} \text{ yr}$, such as Big-Bang nucleosynthesis [3] and anisotropies in the cosmic microwave background radiation [4], yielding upper limits on the long-term averaged variation, $|\dot{G}/G| \lesssim 10^{-13} \text{ yr}^{-1}$. The second class contains measurements on long timescales, $10^9\text{--}10^{10} \text{ yr}$, but without reaching into the very early Universe. In this category, one finds constraints based on paleontology [5] and on stellar astrophysics, the latter including helioseismology [6], binary pulsar masses [7], and globular clusters [8]. Inferred constraints are in the range $|\dot{G}/G| \lesssim 10^{-(11-12)} \text{ yr}^{-1}$. Experiments in the third class measure the change of G directly over short, “human” timescales, $\sim 10 \text{ yr}$. Orbits of planets [9], rotation of isolated pulsars [10], orbits of binary pulsars [11], and oscillations of white dwarfs [12] are some examples of this class, with inferred upper limits $|\dot{G}/G| \lesssim 10^{-(10-12)} \text{ yr}^{-1}$. Even though results from the first category are nominally the most restrictive on a long-term variation of G , they depend on the assumed form of the variation of G near the Big Bang. Thus, it is still useful to consider measurements of the second and third categories, which could detect variations of G in more recent times.

In this paper, we present a new method for constraining a possible time-variation of G , which we call *gravitochemical heating*. A change in G induces a variation in the internal composition of a neutron star, causing dissipation and internal heating. The comparison of the predicted surface temperatures with the only available observation [13] sets constraints on the variation of G . The method is based on the results of Fernández & Reisenegger [14] (see also Refs. [15], [16], and [17]), who showed that internal heating arises as a consequence of the compression of the star due to the decrease of its rotation rate and centrifugal force (*rotochemical heating*). In the present work, compression (or expansion) of the star is caused by an increase (or decrease) in the gravitational force as G changes, with the governing equations having the same mathematical form. Since the chemical relaxation timescale of a neutron star is $\sim 10^8 \text{ yr}$, our experiment is closest to those in the second class discussed above. If G changes over cosmological timescales (as in the proposals of Dirac and Brans-Dicke) $\dot{G}$ can be approximated as a constant, which we will do in most of what follows.

Theory.— The simplest neutron-star models contain only neutrons (n), protons (p), and low-mass leptons (l ; including electrons [e] and muons [μ]), which can convert into each other by beta reactions: direct Urca,

$$\begin{aligned} n &\longrightarrow p + l + \bar{\nu}, \\ p + l &\longrightarrow n + \nu, \end{aligned} \quad (1)$$

possibly forbidden by momentum conservation, and modified Urca,

$$\begin{aligned} n + N &\longrightarrow p + N + l + \bar{\nu}, \\ p + l + N &\longrightarrow n + N + \nu, \end{aligned} \quad (2)$$

where an additional nucleon N must be introduced in order to conserve momentum. Neutrinos (ν) and antineutrinos ($\bar{\nu}$) produced in these reactions escape, taking energy with them and therefore cooling the star [18]. When the latter is in beta equilibrium (with the chemical potentials satisfying $\mu_n = \mu_p + \mu_l$, for both $l = e, \mu$), the rate of direct and inverse reactions is the same, i.e., the composition does not change.

A change in G produces a change in the density of the stellar matter. Since the chemical potentials depend on density, the system departs from the beta equilibrium state. This departure can be quantified by the chemical imbalances,

$$\eta_{npl} = \mu_n - \mu_p - \mu_l. \quad (3)$$

It increases the reaction rates so as to approach a new equilibrium configuration. If G changes continuously, the star is always out of equilibrium, storing an excess of energy that is dissipated as internal heating and enhanced neutrino emission.

The formalism for calculating the evolution of the temperature and chemical imbalances is described in § 2 of Ref. [14]. Here we outline the fundamental equations and the modifications required in order to treat the gravitochemical case. The evolution of the internal temperature as measured by a distant observer, T (we avoid the usual subscript “ ∞ ” in order to keep the equations simple), taken to be uniform inside the star, is given by the thermal balance equation,

$$\dot{T} = \frac{1}{C} (L_H - L_\nu - L_\gamma), \quad (4)$$

where C is the total heat capacity of the star, L_H is the total power released by the heating mechanism (which depends on the chemical imbalances and the temperature), L_ν the total power emitted as neutrinos, and L_γ the power released as thermal photons. The evolution of the chemical imbalances is given by

$$\begin{aligned} \dot{\eta}_{npl} = & -[A_{D,l}(\eta_{npe}, T) + A_{M,l}(\eta_{npe}, T)] \\ & -[B_{D,l}(\eta_{np\mu}, T) + B_{M,l}(\eta_{np\mu}, T)] + C_{npl}\dot{G} \end{aligned} \quad (5)$$

where the subscripts D and M refer to direct and modified Urca reactions, respectively. The functions A and B quantify the effect of reactions towards restoring chemical equilibrium, and thus have the same sign of η_{npl} [14]. The constant C_{npl} quantifies the departure from equilibrium due to $\dot{G}$, being positive and depending on the stellar model and equation of state. It can be further decomposed as $C_{npl} = (Z_{npl} - Z_{np})I_{G,l} + Z_{np}I_{G,p}$, where $I_{G,i} \equiv (\partial N_i^{eq} / \partial G)_A$ is the change of the equilibrium number of each particle species, N_i^{eq} , due to the variation of G , at constant total baryon number A , while Z_i are coefficients depending only on the stellar model (see Ref. [17] for a rigorous calculation of the latter).

As in the rotochemical heating case, the main consequence of gravitochemical heating is that the star arrives at a stationary state, where the rate at which $\dot{G}$ modifies the equilibrium concentrations is the same as the rate at which reactions drive the system toward the new equilibrium configuration, with heating and cooling balancing each other [15]. The properties of this stationary state can be obtained by the simultaneous solution of equations (4) and (5) with $\dot{T} = \dot{\eta}_{npl} = 0$. The existence of the stationary state makes it unnecessary to model the full evolution of the temperature and chemical imbalances of the star in order to calculate the final temperature, since the stationary state is independent of the initial conditions (see Ref. [14] for a detailed analysis of the rotochemical heating case). For a given value of $|\dot{G}|$, it is thus possible to calculate the temperature of an old pulsar that has reached the stationary state, without knowing its exact age.

When only modified Urca reactions operate and nucleon Cooper pairing effects are negligible, it is possible to solve analytically for the stationary values of the photon luminosity L_γ^{st} and chemical imbalances η_{npl}^{st} , as a function of stellar model and current value of $|\dot{G}/G|$. The reason for this is that the longer equilibration timescale given by the slower modified Urca reactions yields stationary chemical imbalances satisfying $\eta_{npl} \gg kT$. The term $L_H - L_\nu$ in the thermal balance equation can be written asymptotically as $K_{Le}\eta_{npe}^8 + K_{L\mu}\eta_{np\mu}^8$, where $K_{L,i}$ are positive constants that depend only on stellar mass and equation of state [14]. Assuming a stellar model with mass $1.41M_\odot$, calculated with the equation of state A18+ δv +UIX* [19], the photon luminosity in the stationary state comes out to be

$$L_\gamma^{st} \simeq 5 \times 10^{28} \left(\frac{|\dot{G}/G|}{10^{-12} \text{ yr}^{-1}} \right)^{8/7} \text{ erg s}^{-1}. \quad (6)$$

Assuming isotropic blackbody radiation, i.e., $L_\gamma = 4\pi\sigma R^2 T_s^4$, with $R \simeq 14$ km the radius of the star as measured by a distant observer, and σ the Stefan-Boltzmann constant, the surface temperature of the star in the stationary state is

$$T_s^{st} \simeq 7.7 \times 10^4 \left(\frac{|\dot{G}/G|}{10^{-12} \text{ yr}^{-1}} \right)^{2/7} \text{ K}. \quad (7)$$

The timescale for the system to reach the stationary state, obtained by the same procedure as in § 4.4 of Ref. [14], is

$$\tau_{st} \simeq 2 \times 10^8 \left(\frac{|\dot{G}/G|}{10^{-12} \text{ yr}^{-1}} \right)^{-6/7} \text{ yr}. \quad (8)$$

Results.— In order to apply this formalism and constrain the value of $|\dot{G}/G|$, we need a neutron star that

(1) has a measured surface temperature (or at least a good enough upper limit on the latter), and (2) is confidently known to be older than the timescale to reach the stationary state for the intended upper limit on $|\dot{G}/G|$. So far, the only object satisfying both conditions is the millisecond pulsar closest to the Solar System, PSR J0437-4715 (hereafter “J0437”), whose surface temperature was inferred from an ultraviolet observation by Kargaltsev, Pavlov, and Romani [13]. Its spin-down age, $\tau_{\text{sd}} \simeq 5 \times 10^9$ yr (e.g., [20]), and the cooling age of its white dwarf companion, $\tau_{\text{WD}} \simeq 2.5 - 5.3 \times 10^9$ yr [22], are much longer than the time required to reach the steady state, as long as $|\dot{G}/G| \gtrsim 10^{-13} \text{ yr}^{-1}$.

Consequently, we consider stellar models constructed from different equations of state, with masses satisfying the constraint obtained for J0437 by van Straten et al. [20], $M_{\text{psr}} = 1.58 \pm 0.18 M_{\odot}$, and calculate the stationary temperature for each, ignoring potential Cooper pairing effects [18]. In Figure 1, we compare the observational constraints on the temperature of this pulsar [13] together with the theoretical prediction for different equations of state (for further details see Table 1 of Ref. [14]), assuming $|\dot{G}/G| = 2 \times 10^{-10} \text{ yr}^{-1}$. When the stellar mass becomes large enough for the central pressure to cross the threshold for direct Urca reactions, T_s drops abruptly, due to the faster relaxation towards chemical equilibrium. This occurs in two steps, as electron and muon direct Urca processes have different threshold densities (see, e.g., Ref. [14]). The chosen value of $\dot{G}$ is such that the stationary temperatures of all stellar models lie (just) above the 90 % confidence contour, as seen in Figure 1, and therefore represents a rather safe, general upper limit.

Nevertheless, conventional neutron star cooling models reproduce observed temperatures better when only modified Urca reactions are considered [18]. Restricting our sample to the equations of state which allow only modified Urca reactions in the mass range considered here, namely A18 + δv , A18 + δv + UIX, BPAL21, and BPAL31, we obtain a more restrictive upper limit, $|\dot{G}/G| < 4 \times 10^{-12} \text{ yr}^{-1}$, illustrated in Figure 2. This value is among the most restrictive results from methods probing a similar timescale.

So far, we have assumed a steady, monotonic time-variation of G . Can our method constrain short-timescale variations? We consider an oscillatory variation, $G(t) = G_0[1 + A \cos(\omega t)]$, in a star with only modified Urca reactions and one leptonic species (electrons), in the interesting limit $\eta_{npe} \gg kT$. The equation for the chemical imbalance evolution is

$$\dot{\eta}_{npe} = -K_A \eta_{npe}^7 + C_{npe} \dot{G}. \quad (9)$$

In the regime of low forcing frequency (when the period is longer than the timescale to reach the steady state), this evolution is equivalent to the situation already considered. In the opposite limit, when the oscillation is




FIG. 1: Comparison of the gravitochemical heating predictions to observations of PSR J0437-4715. The solid lines are the predicted stationary surface temperatures as functions of stellar radius, for different equations of state (A18 from [19] and BPAL from [21]), constrained to the observationally allowed mass range for this pulsar [20]. Dashed lines correspond to the 68% and 90% confidence contours of the blackbody fit of Kargaltsev et al. [13] for the ultraviolet emission from this object. The value of $|\dot{G}/G| = 2 \times 10^{-10} \text{ yr}^{-1}$ is chosen so that all stationary temperature curves lie above the observational constraints. (BPAL32 and BPAL33 allow direct Urca reactions in the observed mass range of J0437.)




FIG. 2: Same as Figure 1, but now the value of $|\dot{G}/G| = 4 \times 10^{-12} \text{ yr}^{-1}$ is chosen so that only the stationary temperature curves with modified Urca reactions are above the observational constraints.

faster than the time required to reach the steady state, we have $\eta_{npe} \simeq C_{npe} A G_0 \cos(\omega t)$. As mentioned above, the photon luminosity satisfies $L_\gamma \simeq K_{Le} \eta_{npe}^8$. Assuming a detection limit similar to the luminosity observed in J0437, $L_\gamma \sim 10^{29} \text{ erg s}^{-1}$, the heating can be detected

only if the oscillation timescale satisfies

$$t_{\text{osc}} \equiv \frac{1}{\omega} \sim \frac{(L_{\gamma}/K_{Le})^{1/8}}{C_{\text{npe}}|\dot{G}|} \gtrsim 3 \times 10^8 \left(\frac{|\dot{G}/G|}{10^{-12} \text{ yr}^{-1}} \right)^{-1} \text{ yr}, \quad (10)$$

too long to be of interest.

Conclusions. – We have shown that a long-term, monotonic variation of G alters the chemical equilibrium state of a neutron star, leading to internal heating. This effect is an exact analog of that produced by spin-down compression [14]. As in that case, the star reaches a stationary state in which the temperature remains nearly constant. We have calculated this temperature as a function of $|\dot{G}/G|$ and stellar mass using realistic equations of state, and compared our results to the temperature inferred from ultraviolet observations of the millisecond pulsar J0437-4715 by Kargaltsev et al. [13]. In the most general case, when direct Urca reactions are allowed to operate, we obtain an upper limit $|\dot{G}/G| < 2 \times 10^{-10} \text{ yr}^{-1}$. Restricting the sample of equations of state to those that allow only modified Urca reactions, we obtain a much more restrictive upper limit, $|\dot{G}/G| < 4 \times 10^{-12} \text{ yr}^{-1}$, competitive with constraints obtained from other methods probing similar timescales. However, since the composition of matter above nuclear densities is uncertain and millisecond pulsars are generally expected to be more massive than classical pulsars, we cannot rule out the result for the direct Urca regime. Information about the prevalence of the direct Urca process and the mass threshold above which it is present may in the future be obtained from the study of the quiescent emission of soft X-ray transients (e. g., Ref. [18]).

Further progress in our knowledge of neutron star matter will allow this method to become more effective at constraining variations in G . The method can also be improved with an increased sample of objects with measured thermal emission. In this respect, an interesting candidate is PSR J0108-1431, plausibly the nearest pulsar, with distance estimates ranging from 60 pc [23] to 184 pc [24]. Due to its much slower rotation (period $P = 0.808 \text{ s}$ [23], compared to $P = 0.00576 \text{ s}$ for J0437 [20]), the current time-derivative of its centrifugal force ($\propto \dot{P}/P^3$) is 680 times smaller than that of J0437. Thus, unless it was born rotating much faster ($P_0 \lesssim 0.02 \text{ s}$), rotochemical heating should be negligible for this pulsar [14, 25], and any detectable surface temperature would be a clear signature of a change in G .

This work was supported by FONDECYT (Chile)

through its Regular Grant 1060644.

-
- [1] P. Dirac, *Nature* (London) **139**, 323 (1937).
 - [2] C. Brans and R. Dicke, *Phys. Rev. Lett.* **124**, 925 (1961).
 - [3] C. J. Copi, A. N. Davis, and L. M. Krauss, *Phys. Rev. Lett.* **92**, 171301 (2004).
 - [4] R. Nagata, T. Chiba, and N. Sugiyama, *Phys. Rev. D* **69**, 083512 (2004).
 - [5] W. Eichendorf and M. Reinhardt, *Mitteilungen der Astronomischen Gesellschaft Hamburg*, **42**, 89 (1977).
 - [6] D. B. Guenther, L. M. Krauss, and P. Demarque, *Astrophys. J.* **498**, 871 (1998).
 - [7] S. E. Thorsett, *Phys. Rev. Lett.* **77**, 1432 (1996).
 - [8] S. Degl’Innocenti *et al.* *Astron. Astrophys.* **312**, 345 (1995).
 - [9] J. G. Williams, X. X. Newhall, and J. O. Dickey, *Phys. Rev. D* **53**, 6730 (1996).
 - [10] I. Goldman, *Mon. Not. R. Astron. Soc.* **244**, 184 (1990).
 - [11] V. M. Kaspi, J. H. Taylor, and M. F. Ryba, *Astrophys. J.* **428**, 713 (1994).
 - [12] O. G. Benvenuto, E. García-Berro, and J. Isern, *Phys. Rev. D* **69**, 082002 (2004).
 - [13] O. Kargaltsev, G. G. Pavlov, and R. Romani, *Astrophys. J.* **602**, 327 (2004).
 - [14] R. Fernández and A. Reisenegger, *Astrophys. J.* **625**, 291 (2005).
 - [15] A. Reisenegger, *Astrophys. J.* **442**, 749 (1995).
 - [16] A. Reisenegger, *Astrophys. J.* **485**, 313 (1997).
 - [17] A. Reisenegger, P. Jofré, R. Fernández, and E. Kantor, *Astrophys. J.*, in press (astro-ph/0606322).
 - [18] D. G. Yakovlev and C. J. Pethick, *Ann. Rev. Astron. Astrophys.* **42**, 169 (2004).
 - [19] A. Akmal, V. R. Pandharipande, and D. G. Ravenhall, *Phys. Rev. C* **58**, 1804 (1998).
 - [20] W. van Straten, M. Bailes, M. Britton, S. R. Kulkarni, S. B. Anderson, R. N. Manchester, and J. Sarkissian, *Nature* (London) **412**, 158 (2001).
 - [21] M. Prakash, T. L. Ainsworth, and J. M. Lattimer, *Phys. Rev. Lett.* **61**, 2518 (1988).
 - [22] B. M. S. Hansen and E. S. Phinney, *Mon. Not. R. Astron. Soc.* **294**, 569 (1998).
 - [23] T. M. Tauris, L. Nicastro, S. Johnston, R. N. Manchester, M. Bailes, A. G. Lyne, J. Glowacki, D. R. Lorimer, and N. D’Amico, *Astrophys. J.* **428**, L53 (1994).
 - [24] Calculated online from the Cordes-Lazio NE2001 Galactic free electron distribution model, http://rsd-www.nrl.navy.mil/7213/lazio/ne_model/ (J. M. Cordes and T. J. W. Lazio, preprint astro-ph/0207156).
 - [25] D. González, A. Reisenegger, and R. Fernández, paper in preparation (2006).